

# KSA Algae Living Library

Microalgae and cyanobacteria isolated from the Kingdom of Saudi Arabia

Dr. Bárbara Bastos de Freitas Prof. Kyle J. Lauersen







Biotechnology group







# KSA Algae Living Library – a resource of locally isolated microalgae and cyanobacteria from the Kingdom of Saudi Arabia

This document contains information on microalgae and cyanobacteria which have been isolated from numerous environments within the Kingdom of Saudi Arabia (KSA) that are part of the Living Library of KSA species maintained within the Sustainable & Synthetic Biotechnology (SSB) group on KAUST campus. Information found in this document includes, where available, the species identifiers, microscopy image(s), sample location, as well as preliminary growth data when available. New species are routinely being isolated and added to this collection. Those featured here without full definitions or growth information are in process and will be added to this living document as data becomes available.

The main goal of this project is isolate, cultivate, characterize and generate a living library of KSA photosynthetic microbes that can be cultivated in salt or fresh waters, for carbon, nitrogen, and phosphorous waste-stream reuse applications. This work aims to support Vision2030 in alignment with the The Research, Development and Innovation Authority (RDIA) pillars.

**Authors:** Dr. Bárbara Catarina Bastos de Freitas has led isolation efforts and project management in the lab of Prof. Kyle J. Lauersen.

**Technical support**: Activities in the SSB Lab are currently supported by Mirian dos Santos Mendes. Previous support was provided by Dr. Adriana Barsotti. Bioinformatics support is supported by Mark Pampuch.

# Direct financial support for this effort has been provided by:

- KAUST Circular Carbon Initiative, grants 4769 & 5213 "Bioprospecting algal species from the Red Sea and KSA for carbon reuse and conversion to commodities" & "Scaling production of high value biochar: feedstock variability, material characterization, and field trials"
- KAUST Center of Excellence in Sustainable Food Security, grant 5934 "Demonstration facility for sustainable high-value proteins from low-value waste."
- KAUST Baseline Research Funding awarded to Prof. Kyle J. Lauersen

Collaborative support: From 2022-2023, some strain genetic and metabolite analyses which do not appear in this document were conducted by Dr. Cristina Andres-Barrao and Dr. Rahul Kapoor, and also knowledge exchange about isolation techniques was shared from Mr. Gabriel Renato Castro, from the Development of Algae Biotechnology in Kingdom of Saudi Arabia (DAB-KSA) project which is a Ministry of Environment Water and Agriculture (MEWA) funded research project managed by KAUST Beacon Development. Other assistance with sample collection at different locations is noted on individual isolate pages.

This document was last updated on: November 02, 2025

# **Table of Contents**

1	Introduction	1
2	Notes	2
3	Microalgae within the Living Library	4
	KAUST 001 - Synechococcus elongatus	
	KAUST 002 - Synechococcus elongatus Strain Identification	
	KAUST 003 - Trebouxiophyceae sp	
	KAUST 004 - Parachlorella kessleri Strain Identification	
	KAUST 005 - Tetradesmus obliquus Strain Identification	
	KAUST 006 - Chlorella sp	
	KAUST 007 - Auxenochlorella pyrenoidosa Strain Identification	
	KAUST 008 - Asterarcys sp	
	KAUST 009 - Enallax costatusStrain Identification	
	KAUST 010 - Monoraphidium sp	
	KAUST 011 - Chlorella sp	
	KAUST 012 - Synechococcus elongatus Strain Identification	
	KAUST 013 - Chlorella sorokiniana Strain Identification	
	KAUST 014 - Chlorella sp	
	KAUST 015 – Chlorella sp	
	KAUST 016 - Micractinium sp	21
	KAUST 017 - Coelastrella sp	22
	KAUST 018 - Chlorella thermophila Strain Identification	23
		23

24
24
25 25
26
27 27
28 28
29
3 <i>c</i>
31
32 32
<i>33</i>
34
35
36
37
38 38
39
40 40
41
42
43
44

KAUST 040 – "CCAP 293/7"Strain Identification	
KAUST 041 – "CCAP 293/9" Strain Identification	46
KAUST 042 – "CCAP 293/8" Strain Identification	47
KAUST 043 – "CCAP 293/10" Strain Identification	
KAUST 044 - "CCAP 1479/18" Strain Identification	
KAUST 045 – "CCAP 1479/22" Strain Identification	
KAUST 046 – Chlorella sorokiniana Strain Identification	
KAUST 047 - Green alga of currently unknown identity Strain Identification	
KAUST 048 - Chlamydomonas reinhardtii Strain Identification	
KAUST 049 - Trichocoleus desertorum Strain Identification	
KAUST 050 - Green alga of currently unknown identity Strain Identification	
KAUST 051 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 052 - Green alga of currently unknown identity Strain Identification	
KAUST 053 - Chlamydomonadales sp Strain Identification	
KAUST 054 - Dunaliella polymorpha Strain Identification	
KAUST 055 - Lobochlamys sp Strain Identification	
KAUST 056 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 057 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 058 - Diatom of currently unknown identity Strain Identification	
KAUST 059 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 060 - Cyanobacterium of currently unknown identity	

KAUST 061 - Pleodorina illinoisensis	
KAUST 062 - Coelastrum microporumStrain Identification	
KAUST 063 - Monoraphidium circinale	
KAUST 064 - Chlamydomonas sp	
KAUST 065 – Phormidesmis molle	
KAUST 066 – Desmodesmus sp. Strain Identification	
KAUST 067 - Scenedesmus spStrain Identification	
KAUST 068 - Nitzschia spStrain Identification	
KAUST 069 - Anabaenopsis sp. Strain Identification	
KAUST 070 - Fistulifera saprophila	
KAUST 071 – Desmodesmus sp	
KAUST 072 - Chlamydomonas sp	
KAUST 073 - Green alga of currently unknown identity	
KAUST 074 - Tetranephris brasiliensis	
KAUST 075 - Green alga of currently unknown identity	
KAUST 076 - Green alga of currently unknown identity	81
KAUST 077 - Stichococcus mirabilis	82
KAUST 078 - Chlamydomonas reinhardtii	83
KAUST 079 - Diatom of currently unknown identity	84
KAUST 080 - Diatom of currently unknown identity	85
KAUST 081 - Green alga of currently unknown identity	86

KAUST 082 - Green alga of currently unknown identity  Strain Identification	
KAUST 083 - Nitzschia sp.	
Strain Identification	
KAUST 084 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 085 - Cyanobacterium of currently unknown identity Strain Identification	90
KAUST 086 - Green alga of currently unknown identity Strain Identification	
KAUST 087 - Chlamydomonas isabeliensis Strain Identification	
KAUST 088 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 089 - Green alga of currently unknown identity Strain Identification	
KAUST 090 – Synechocystis sp Strain Identification	
KAUST 091 - Green alga of currently unknown identity Strain Identification	
KAUST 092 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 093 – Cyanobacterium of currently unknown identity Strain Identification	
KAUST 094 - Heterochlorella luteoviridis	
KAUST 095 - Chlamydomonas sp. Strain Identification	
KAUST 096 - Green alga of currently unknown identity Strain Identification	
KAUST 097 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 098 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 099 - Green alga of currently unknown identity Strain Identification	
KAUST 100 - Viridiparva madeirensis Strain Identification	
KAUST 101 - Green alga of currently unknown identity Strain Identification	
KAUST 102 - Stichococcus sp	

KAUST 103 - Cyanobacterium of currently unknown identity	
KAUST 104 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 105 - Green alga of currently unknown identity Strain Identification	
KAUST 106 - Green alga of currently unknown identity Strain Identification	
KAUST 107 - Green alga of currently unknown identity Strain Identification	
KAUST 108 - Cyanobacterium of currently unknown identity Strain Identification	
KAUST 109 - Green alga of currently unknown identity Strain Identification	
KAUST 110 - Green alga of currently unknown identity Strain Identification	
KAUST 111 – Chlorella sorokiniana Strain Identification	
KAUST 112 – Parachlorella kessleri Strain Identification	
KAUST 113 – Chlorella vulgaris Strain Identification	
KAUST 114 – Chlorella sp	
KAUST 115 – Chlorella sorokiniana Strain Identification	
KAUST 116 – Micractinium sp	
KAUST 117 – Chlorella sp	
KAUST 118 – Leptolyngbya breviarticulata Strain Identification	
KAUST 119 – Green alga of currently unknown identity Strain Identification	
KAUST 120 – Cyanobacterium of currently unknown identity Strain Identification	
KAUST 121 – Diatiom of currently unknown identity Strain Identification	
KAUST 122 – Green alga of currently unknown identity	
KAUST 123 — Cyanobacterium of currently unknown identity	

KAUST 124 - Diatom of currently unknown identity	
Strain Identification	129
KAUST 125 – Diatom of currently unknown identity Strain Identification	
KAUST 126 – Cyanobacterium of currently unknown identity Strain Identification	
KAUST 127 – Cyanobacterium of currently unknown identity Strain Identification	
KAUST 128 – Green alga of currently unknown identity Strain Identification	
KAUST 129 – Diatom of currently unknown identity Strain Identification	
KAUST 130 – Asterarcys sp	
KAUST 131 – Chlamydomonas sp	
KAUST 132 – Chlorella sp	
KAUST 133 – Tetraselmis sp	
KAUST 134 – Chlorella sorokinina	
KAUST 135 – Dunaliella tertiolecta	
KAUST 136 — Monoraphidium subclavatumStrain Identification	
KAUST 137 – Diatom of currently unknown identity	
KAUST 138 — Chlamydomonas sp	
KAUST 139 – Porphyridium sp	
KAUST 140 - Cyanobacterium of currently unknown identity	
KAUST 141 — Cyanobacterium of currently unknown identity	
KAUST 142 – Green alga of currently unknown identity	
KAUST 143 — Cyanobacterium of currently unknown identity	
KAUST 144 – Cyanobacterium of currently unknown identity	

KAUST 145 – Cyanobacterium of currently unknown identity	150
Strain Identification	
KAUST 146 – Cyanobacterium of currently unknown identity	
Strain Identification	151
KAUST 147 – Scenedesmus sp	
	_
KAUST 148 – Synechocystis sp Strain Identification	
KAUST 149 – Rhizoclonium sp	154
Strain Identification	
KAUST 150 – Cyanobacterium of currently unknown identity	155
Strain Identification	
KAUST 151 – Cyanobacterium of currently unknown identity	156
Strain Identification	
KAUST 152 - Cyanobacterium of currently unknown identity	157
Strain Identification	
KAUST 153 - Cyanobacterium of currently unknown identity	158
Strain Identification	
KAUST 154 - Diatom of currently unknown identity	159
Strain Identification	
KAUST 155 - Green alga of currently unknown identity	160
Strain Identification	
KAUST 156 - Green alga of currently unknown identity	161
Strain Identification	

#### 1 Introduction

Algae are diverse organisms which can be found in all environments around the globe. They are ancient and have complex evolutionary histories, most of these organisms are photosynthetic, using light for energy and carbon dioxide (CO<sub>2</sub>) as a carbon source for growth. Like plants, algae naturally represent carbon reuse vehicles as they convert inorganic CO<sub>2</sub> into all of their respective bio-molecules: proteins, lipids, carbohydrates, and specialty chemicals but they do not produce the complex structural tissues of their higher-plant counterparts.

Algae can be grown with simple inputs such as trace minerals, carbon dioxide, (sun)light, and water, with some single-celled microalgae exhibiting rapid doubling times that far exceed the growth rates of higher plants. Their cultivation can be performed in photobioreactors, built on any land-type, use wasteor sea water, do not compete with farming resources, and can even be coupled to point sources of CO<sub>2</sub> emissions.

Algae are polyphyletic, and have evolved different properties related to their individual environmental niches. This biodiversity is barely explored and requires increased attention as each may hold unique value for different human-use applications, indeed algal biomass itself has been developed into a valuable commodity in the food and medicinal sectors. Different algae have specialized value for different reasons, some hyper accumulate triacylglycerols (oils), which can be converted into hydrocarbon or oleochemicals, others secrete polysaccharides, which have value as medicines and cosmetics, others produce specialty pigments, or even specialty antiviral compounds. As the Kingdom of Saudi Arabia is looking to diversify its sustainable technologies, the development and bio-prospecting of local algae strains from the Kingdom may hold value for a number of industries. Sourcing local strains is of importance for their ability to cope with the harsh environments of this region in outdoor cultivation concepts and to avoid issues of invasive species or cross border intellectual property with regards to the Nagoya protocol. The extreme environments of the region, have likely yielded microbes with advanced biochemistries, capable of surviving the harsh UV and light radiation as well as high temperatures. Algal biodiversity has been generally underexplored in the KSA.

#### 2 Notes

We performed morphological identification of the isolated algae by microscopic observation. The strains were molecularly identified by amplification and sequencing using different primers (listed in Table 1). The total DNA was extracted from frozen cellular pellets (pellets were obtained by centrifuging 2 mL of each monoculture) using Quick-DNA Fungal/Bacterial Miniprep Kit (Zymo Research). The PCR reactions were performed using Phusion Green Hot Start II High-Fidelity PCR Master Mix (Thermo Scientific). PCR clean-up was performed with DNA Clean. & Concentrator<sup>TM</sup> -25 (Zymo Research), and Sanger sequencing was performed on PCR products.

Other data to be collected includes growth data and biomass composition. We tested the feasibility of cultivating the local strains using outdoor light and temperature conditions of Thuwal, Saudi Arabia (22°18'16.6"N 39°06'07.9"E) by weather simulation in Algem photobioreactors. For each local strain, we grew six experimental conditions: constant light, 12 h:12 h light:dark, either at 25°C or 30°C and 325  $\mu$ mol photons m<sup>-2</sup> s<sup>-1</sup> ( $\mu$ E) as controls. and 4 different temperature and light conditions that each represent one of the four seasons in Thuwal, Saudi Arabia. The weather data used for this experiment consisted of temperature, and photosynthetically active radiation (PAR) measurements (10 min intervals) recorded at the Coastal and Marine Resources Core Lab (CMR) at KAUST between January and December 2014. Local weather station data were used to generate a representative week from each of February (Winter), May (Spring), August (Summer), and November (Autumn) from 2014 in the reactors. This is a living document and will be updated regularly.


For inquiries regarding growth data, biomass composition, and comprehensive details on growth conditions, we invite you to make formal inquiries. Requests for this information can be directed to Professor Kyle J. Lauersen and Dr. Bárbara Bastos de Freitas (Curator), who will provide guidance on the protocol for processing such requests.

Primer	Marker (locus/gene;	Sequence	Annealing	Designed	Product
	abbreviated name		temperature	specificity	size
			(°C)		(bp)
Cdm F	18S ribosomal DNA;	GTCAGAGGTGAAATTCTTGGATTTA	62.2	Universal	730
Cdm R	small subunit; 18S	AAGGGCAGGGACGTAATCAACG	67.7	algae	
ITS1 F modified	Internal transcribed	AGGAGAAGTCGTAACAAGGT	61.3	Universal	250-
ITS2T R	spacer 1; ITS1	TTCGCTGCGTTCTTCATCGTT	65.9	algae	1000
Cya106 F	16S ribosomal DNA;	CGGACGGGTGAGTAACGCGTGA	71.2	Cyanobacteria	500
Cya781b R	Cya16S	GACTACAGGGGTATCTAATCCCTTT	64.2		
RbcL F	Ribulose-1,5-	AACCTTTCATGCGTTGGAGAGA	65.3	Universal	500
RbcL R	bisphosphate	CCTGCATGAATACCACCAGAAGC	66.5	algae	
	carboxylase/oxygenase				
	-Large subunit; rbcL				
	, , , , ,				

# 3 Microalgae within the Living Library

The following pages contain information sheets on each isolate, the strain code contains a number which represents its sample.

Location data of sampling is provided in longitude and latitude. Whenever possible a picture of the site is provided. GIS visualization by Abhishekh P. Vijayan (Figure 1). The sampling collection location is shown in Figure 2 and will be updated regularly.



**Figure 1.** GIS visualization.




**Figure 2.** Sample collection location (the green markers represent the location where sampling was conducted).

Identification of new isolates follows the following steps

- Isolation to monoculture
- Microscopic morphological identification
- Genetic identification and assessment of species identity

# **KAUST 001 - Synechococcus elongatus**





Freshwater *Synechococcus elongatus* isolated from a pond in a hot spring outside of town Al Lith, located southwest of Mecca in the Red Sea coast region.

Cultivation medium: BG-11


GPS coordinate: 20°27'35.3"N 40°28'11.0"E

# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 16S ribosomal DNA – Cya106 F/Cya781b R- used in a BLAST search, the sequence identity of 100% shows that *Synechococcus elongatus* PCC 11802 is a relative.

#### **KAUST 002 - Synechococcus elongatus**





Freshwater *Synechococcus elongatus* isolated from the mud in a hot spring outside of town Al Lith, located southwest of Mecca in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 20°27'35.3"N 40°28'11.0"E

#### Strain Identification

Besides the morphological taxonomy, taxonomic classification was evaluated by first performing a genome assembly of the strain, followed by 16s rRNA sequence isolation using Barrnap and a k-mer based classification using Kraken2. rRNA sequences were searched against the NCBI database using BLAST. Both analyses show "Synechococcus elongatus" is a relative, with a BLAST results showing a sequence identity of 99.93%.

#### KAUST 003 - Trebouxiophyceae sp.





Freshwater *Trebouxiophyceae* sp. isolated from a hot spring outside of town Al Lith, located southwest of Mecca in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 20°27'35.3"N 40°28'11.0"E

# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Trebouxiophyceae* sp. BZ-2019 is a relative.

#### KAUST 004 - Parachlorella kessleri





Freshwater *Parachlorella kessleri* isolated from a hot spring outside of town Al Lith, located southwest of Mecca in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 20°27'35.3"N 40°28'11.0"E

# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 100% shows that *Parachlorella kessleri* is a relative.

#### **KAUST 005 - Tetradesmus obliquus**





Freshwater *Tetradesmus obliquus* isolated from a hot spring outside of town Al Lith, located southwest of Mecca in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 20°27'35.3"N 40°28'11.0"E

# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 99.60% shows that *Tetradesmus obliquus* is a relative.

\*This strain was shared with the DAB-KSA team in February/2023 and August/2024 for further analysis in scale-up cultivations.

#### KAUST 006 - Chlorella sp.





Freshwater *Chlorella* sp. KU211 isolated from a hot spring outside of town Al Lith, located southwest of Mecca in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 20°27'35.3"N 40°28'11.0"E


# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 100% shows that *Chlorella* sp. KU211 is a relative.

<sup>\*</sup>This strain was shared with the DAB-KSA team in August/2024 for further analysis in scale-up cultivations.

#### KAUST 007 - Auxenochlorella pyrenoidosa





Seawater Auxenochlorella pyrenoidosa isolated from a hot spring outside of town Al Lith, located southwest of Mecca in the Red Sea coast region. Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 20°27'35.3"N 40°28'11.0"E

# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 100% shows that *Auxenochlorella pyrenoidosa* is a relative.

#### KAUST 008 - Asterarcys sp.





Freshwater *Asterarcys* sp. isolated at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.

Cultivation medium: BG-11


GPS coordinate: 22°18'28.0"N 39°06'23.0"E

# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase -Large subunit -RbcL F /RbcL R - used in a BLAST search, the sequence identity of 99.65% shows that *Asterarcys* sp. is a relative.

#### **KAUST 009 - Enallax costatus**





Freshwater *Enallax costatus* isolated at King Abdullah University of Science and Technology (Thuwal, Makkah), in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 22°18'28.0"N 39°06'23.0"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase -Large subunit -RbcL F /RbcL R - used in a BLAST search, the sequence identity of 98.46% shows that *Enallax costatus* is a relative.

#### KAUST 010 - Monoraphidium sp.





Freshwater *Monoraphidium* sp. at King Abdullah University of Science and Technology (Thuwal, Makkah), in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 22°18'28.0"N 39°06'23.0"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 100% shows that *Monoraphidium* sp. Zaqeq 32 is a relative.

\*This strain was shared with the DAB-KSA team in August/2024 for further analysis in scale-up cultivations.

#### KAUST 011 - Chlorella sp.





Freshwater *Chlorella* sp. isolated from the desert region of Al Ula, located in the Medina Region in north-western Saudi Arabia.

Cultivation medium: BG11

GPS coordinate: 26°36'59.99"N 37°54'59.99"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase -Large subunit -RbcL F /RbcL R - used in a BLAST search, the sequence identity of 100% shows that *Chlorella sp.* MOW 10 is a relative.

\*This strain was shared with the DAB-KSA team in August/2024 for further analysis in scale-up cultivations.

#### **KAUST 012 - Synechococcus elongatus**





Freshwater *Synechococcus elongatus* isolated from the Deserts regions of Al Ula, located in the Medina Region in north-western Saudi Arabia.

Cultivation medium: BG-11

GPS coordinate: 26°36'59.99"N 37°54'59.99"E

#### Strain Identification

Besides the morphological taxonomy, taxonomic classification was evaluated by first performing a genome assembly of the strain, followed by 16s rRNA sequence isolation using Barrnap and a k-mer based classification using Kraken2. rRNA sequences were searched against the NCBI database using BLAST. Both analyses show "Synechococcus elongatus" is a relative, with a BLAST results showing a sequence identity of 99.93%.

#### KAUST 013 - Chlorella sorokiniana





Freshwater *Chlorella sorokiniana* isolated from a hot spring called Al Lith, located southwest of Mecca in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 20°27'35.3"N 40°28'11.0"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Chlorella sorokiniana* is a relative.

#### KAUST 014 - Chlorella sp.





Freshwater *Chlorella* sp. isolated from the Deserts regions of Al Ula, located in the Medina Region in north-western Saudi Arabia.

Cultivation medium: BG-11

GPS coordinate: 26°36'59.99"N 37°54'59.99"E

# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 99.69% shows that *Chlorella* sp. is a relative.

\*This strain was shared with the DAB-KSA team in August/2024 for further analysis in scale-up cultivations.

#### KAUST 015 - Chlorella sp.





Freshwater *Chlorella* sp. isolated from the Deserts regions of Al Ula, located in the Medina Region in north-western Saudi Arabia.

Cultivation medium: BG-11

GPS coordinate: 26°36'59.99"N 37°54'59.99"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase -Large subunit -RbcL F /RbcL R - used in a BLAST search, the sequence identity of 100% shows that *Chlorella sp.* MOW 12 is a relative.

#### KAUST 016 - Micractinium sp.





Freshwater *Micractinium* sp. isolated from a hot spring called Al Lith, located southwest of Mecca in the Red Sea coast region.


Cultivation medium: BG-11

GPS coordinate: 20°27'35.3"N 40°28'11.0"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Micractinium* sp. is a relative.

#### KAUST 017 - Coelastrella sp.





Freshwater *Coelastrella* sp. CORE isolated from Anaerobic Membrane Bioreactor (AnMBr) effluent at King Abdullah University of Science and Technology (Thuwal, Makkah), in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 22°18'35.8"N 39°06'12.4"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 100% shows that *Coelastrella* sp. CORE -2 is a relative.

#### KAUST 018 - Chlorella thermophila





Freshwater *Chlorella thermophila* isolated from a hot spring called Al Lith, located southwest of Mecca in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 20°27'35.3"N 40°28'11.0"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 99.42% shows that *Chlorella thermophila* is a relative.

# KAUST 019 - Raphidocelis subcapitata





Freshwater *Raphidocelis subcapitata* isolated from a greenhouse at National Aquaculture Group (NAQUA) facilities.

Cultivation medium: BG-11

GPS coordinate: 20°19'16.1"N 40°02'27.9"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 96.09% shows that *Raphidocelis subcapitata* is a relative.

\*This strain resulted from environmental samples taken on a field trip to National Aquaculture Group (NAQUA) facilities organized by the Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA).

# KAUST 020 - Monoraphidium dybowskii





Freshwater *Monoraphidium dybowskii* isolated from a greenhouse at National Aquaculture Group (NAQUA) facilities.

Cultivation medium: BG-11

GPS coordinate: 20°19'16.1"N 40°02'27.9"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 99.26% shows *Monoraphidium dybowskii* is a relative.

\*This strain resulted from environmental samples taken on a field trip to National Aquaculture Group (NAQUA) facilities organized by the Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA).

\*This strain was shared with the DAB-KSA team in February/2023 and August/2024 for further analysis in scale-up cultivations.

#### KAUST 021 - Asterarcys sp.





Freshwater *Asterarcys* sp. isolated from a greenhouse at National Aquaculture Group (NAQUA) facilities.

Cultivation medium: BG-11

GPS coordinate: 20°19'16.1"N 40°02'27.9"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase -Large subunit -RbcL F /RbcL R - used in a BLAST search, the sequence identity of 99.70% shows *Asterarcys* sp. GP-2019 is a relative.

\*This strain resulted from environmental samples taken on a field trip to National Aquaculture Group (NAQUA) facilities organized by the Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA).

#### KAUST 022 - Nostoc sp.





Freshwater *Nostoc* sp. from a greenhouse at National Aquaculture Group (NAQUA) facilities.

Cultivation medium: BG-11

GPS coordinate: 20°19'16.1"N 40°02'27.9"E


### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 16S ribosomal DNA – Cya106 F/Cya781b R- used in a BLAST search, the sequence identity of 100% shows that *Nostoc* sp. PCC 7423 is a relative.

\*This strain resulted from environmental samples taken on a field trip to National Aquaculture Group (NAQUA) facilities organized by the Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA).

### KAUST 023 - Leptolyngbya sp.





Freshwater *Leptolyngbya* sp. isolated from a greenhouse at National Aquaculture Group (NAQUA) facilities.

Cultivation medium: BG-11

GPS coordinate: 20°19'16.1"N 40°02'27.9"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 16S ribosomal DNA – Cya106 F/Cya781b R- used in a BLAST search, the sequence identity of 100% shows that *Leptolyngbya* sp. CENA131 is a relative.

\*This strain resulted from environmental samples taken on a field trip to National Aquaculture Group (NAQUA) facilities organized by the Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA).

\*This strain was shared with the DAB-KSA team in August/2024 for further analysis in scale-up cultivations.

### **KAUST 024 - Diatom of currently unknown identity**





Seawater diatom isolated from a pond at National Aquaculture Group (NAQUA) facilities.

Cultivation medium: Red Sea water medium with f/2 nutrients.


GPS coordinate: 20°19'17.4"N 40°02'27.9"E

### Strain Identification

\*This strain resulted from environmental samples taken on a field trip to National Aquaculture Group (NAQUA) facilities organized by the Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA).

### KAUST 025 - Chlorokybus sp.





Freshwater green alga isolated from a biocrust in a volcano area (Harrat Kishb - lava field).

Cultivation medium: BG-11


GPS coordinate: 22°54'00.0"N 41°08'12.1"E

Strain Identification

\*This strain resulted from environmental samples taken on a field trip to National Aquaculture Group (NAQUA) facilities organized by the Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA).

#### **KAUST 026 - Chlorella variabilis**





Freshwater *Chlorella variabilis* isolated from the soil of a greenhouse at National Aquaculture Group (NAQUA) facilities.

Cultivation medium: BG-11

GPS coordinate: 20°16'13.0"N 39°55'46.0"E


### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase-Large subunit-RbcL F/RbcL R - used in a BLAST search, the sequence identity of 100% shows that *Chlorella variabilis* is a relative.

\*This strain resulted from environmental samples taken on a field trip to National Aquaculture Group (NAQUA) facilities organized by the Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA).

### KAUST 027 - Chlamydomonas sp.





Freshwater "Chlamydomonas sp." isolated from soil samples from Al Wahba Crater.

Cultivation medium: BG-11

GPS coordinate: 22°54'22.0"N 41°08'07.0"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 92.71% shows that the strain currently identified as "Chlamydomonas sp. HyS1201A1" is a relative.

\*Chlamydomonas taxonomy is currently undergo138ing revision, and true identity by ITS is not conclusive.

Therefore, the name is indicated in quotations.

### KAUST 028 - Cyanobacterium aponinum





Freshwater *Cyanonbacterium aponinum* isolated from AlSafwa cement company.

Cultivation medium: BG-11

GPS coordinate: 22°33'17.0"N 39°26'11.0"E

### Strain Identification

Besides the morphological taxonomy, taxonomic classification was evaluated by first performing a genome assembly of the strain, followed by 16s rRNA sequence isolation using Barrnap and a k-mer based classification using Kraken2. rRNA sequences were searched against the NCBI database using BLAST. Both analyses show "Cyanobacterium aponium" is a relative, with a BLAST results showing a sequence identity of 100%.

# KAUST 029 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from AlSafwa cement company. Cultivation medium: BG-11

GPS coordinate: 22°33'17.0"N 39°26'11.0"E

Strain Identification

**KAUST 030 - Diatom of currently unknown identity** 





Seawater diatom isolated from Red Sea in the coral region called Ana Coral.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 21°39'19.7"N 38°52'27.1"E

# Strain Identification

### KAUST 031 - *Ulvophyceae* sp.





Seawater *Ulvophyceae* sp. isolated from Red Sea in the coral region called Ana Coral.

Cultivation medium: Red Sea water medium with f/2 nutrients.


GPS coordinate: 21°39'19.7"N 38°52'27.1"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 96.09% shows that *Ulvophyceae* sp. is a relative.

### KAUST 032 - Ulvophyceae sp.





Seawater *Ulvophyceae* sp. isolated from Red Sea in the coral region called Rose Reef.
Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 21°39'19.7"N 38°52'27.1"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Ulvophyceae* sp. *MBIC10479* is a relative.

### KAUST 033 – Leptolyngbya tenuis





Seawater *Leptolyngbya tenuis* isolated from Red Sea in the coral region called Cement Reef.
Cultivation medium: Red Sea water medium with f/2 nutrients.


GPS coordinate: 22°22'26.0"N 38°51'20.0"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 16S ribosomal DNA – Cya106 F/Cya781b R- used in a BLAST search, the sequence identity of 100% shows that *Leptolyngbya tenuis* is a relative.

KAUST 034 - Green alga of currently unknown identity





Seawater green alga isolated from Red Sea in the coral region called Cement Reef.


Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°22'26.0"N 38°51'20.0"E

Strain Identification

# KAUST 035 - Green alga of currently unknown identity





Freshwater green alga isolated from volcanic soil (Harrat Kishb - lava field).

Cultivation medium: BG-11

GPS coordinate: 22°54'00.0"N 41°08'12.1"E

Strain Identification

### **KAUST 036 - Chromochloris zofingiensis**





Freshwater *Chromochloris zofingiensis* isolated from a biocrust in Taif.

Cultivation medium: BG-11

GPS coordinate: 21°17'02.8"N 40°25'29.3"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 96.69% shows that *Chromochloris zofingiensis* is a relative.

### **KAUST 037 - Nostoc commune**





Freshwater *Nostoc commune* isolated from a biocrust in Taif.

Cultivation medium: BG-11

GPS coordinate: 21°17'02.8"N 40°25'29.3"E

Strain Identification

# KAUST 038 - Green alga of currently unknown identity





Freshwater green alga isolated from a white volcano area (Harrat Khaybar- Felsic volcano).


Cultivation medium: BG-11

GPS coordinate: 25°35'18.6"N 39°45'25.1"E

Strain Identification

### KAUST 039 - Dunaliella sp.





Seawater *Dunaliella* isolated from a pond at National Aquaculture Group (NAQUA) facilities. Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 20°19'07.7"N 40°02'44.9"E

Strain Identification

\*This strain resulted from environmental samples taken on a field trip to National Aquaculture Group (NAQUA) facilities organized by the Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA).

# KAUST 040 - "CCAP 293/7"




Unicellular Microulva isolated by John Archer's team.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°22'37.8"N 39°03'18.1"E

Strain Identification

# KAUST 041 - "CCAP 293/9"



Unicellular Microulva isolated by John Archer's team.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°22'37.8"N 39°03'18.1"E

# Strain Identification

# KAUST 042 - "CCAP 293/8"



Unicellular Microulva isolated by John Archer's team.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°22'37.8"N 39°03'18.1"E

# Strain Identification

### KAUST 043 - "CCAP 293/10"




Unicellular Microulva isolated by John Archer's team.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°22'37.8"N 39°03'18.1"E

Strain Identification

# KAUST 044 - "CCAP 1479/18"




Unicellular Microulva isolated by John Archer's team.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°22'37.8"N 39°03'18.1"E

Strain Identification

# KAUST 045 - "CCAP 1479/22"



Unicellular Microulva isolated by John Archer's team.


Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°22'37.8"N 39°03'18.1"E

Strain Identification

#### KAUST 046 – Chlorella sorokiniana





Freshwater *Chlorella sorokiniana* isolated at King Abdullah University of Science and Technology (Thuwal, Makkah), in the Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 22°18'29.0"N 39°06'23.0"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Chlorella sorokiniana isolate LWG002615* is a relative.

# KAUST 047 - Green alga of currently unknown identity





Freshwater green alga isolated at a volcanic area

– Al Wahbah well.


Cultivation medium: BG-11

GPS coordinate: 22°54'22.0"N 41°08'07.0"E

Strain Identification

### KAUST 048 - Chlamydomonas reinhardtii





Freshwater "Chlamydomonas reinhardtii" isolated from an Oasis at a volcanic area – Al Wahbah well.

Cultivation medium: BG-11

GPS coordinate: 22°54'22.0"N 41°08'07.0"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 86% shows that "Chlamydomonas reinhardtii" is a relative.

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity by ITS is not conclusive. Therefore, the name is indicated in quotations.

#### KAUST 049 - Trichocoleus desertorum





Freshwater *Trichocoleus desertorum* isolated at Moon Valley desert.

Cultivation medium: BG-11

GPS coordinate: 23°37'36.8"N 38°48'29.0"E

### Strain Identification

Besides the morphological taxonomy, taxonomic classification was evaluated by first performing a genome assembly of the strain, followed by 16s rRNA sequence isolation using Barrnap and a k-mer based classification using Kraken2. rRNA sequences were searched against the NCBI database using BLAST. Both analyses show "*Trichocoleus desertorum*" is a relative, with a BLAST results showing a sequence identity of 93.64%.

# KAUST 050 - Green alga of currently unknown identity





Freshwater green alga isolated at Moon Valley desert.

Cultivation medium: BG-11

GPS coordinate: 23°37'36.8"N 38°48'29.0"E

# Strain Identification

KAUST 051 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from tanks in the city of Asfan.


Cultivation medium: BG-11

GPS coordinate: 21°50'34.0"N 39°24'14.6"E

Strain Identification

# KAUST 052 - Green alga of currently unknown identity





Freshwater green alga isolated from tanks in the city of Asfan.

Cultivation medium: BG-11

GPS coordinate: 21°50'34.0"N 39°24'14.6"E

Strain Identification

### KAUST 053 - Chlamydomonadales sp.





Seawater "Chlamydomonadales sp." isolated from Al Wahbah crater.

Cultivation medium: Red Sea water medium with f/2 nutrients..

GPS coordinate: 22°54'04.0"N 41°08'21.1"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 97.59% shows that "Chlamydomonadales sp." is a relative.

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity by 18S is not conclusive. Therefore, the name is indicated in quotations.

### KAUST 054 - Dunaliella polymorpha





Seawater *Dunaliella polymorpha* isolated from Al Wahbah crater.

Cultivation medium: Red Sea water medium with f/2 nutrients..


GPS coordinate: 22°54'04.0"N 41°08'21.1"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 77.87% shows that *Dunaliella polymorpha* is a relative.

### KAUST 055 - Lobochlamys sp.





Seawater "Lobochlamys sp." isolated from Al Wahbah crater.

Cultivation medium: Red Sea water medium with f/2 nutrients..

GPS coordinate: 22°54'04.0"N 41°08'21.1"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 97% shows that "Lobochlamys sp." is a relative.

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity by 18S is not conclusive. Therefore, the name is indicated in quotations.

# KAUST 056 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from Al Wahbah crater.

Cultivation medium: BG-11.

GPS coordinate: 22°54'04.0"N 41°08'21.1"E

Strain Identification

### KAUST 057 - Cyanobacterium of currently unknown identity





Seawater cyanobacterium isolated from North Beach at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.

Cultivation medium: Red Sea water medium with f/2 nutrients..

GPS coordinate: 23°37'02.6"N 38°48'17.8"E

### Strain Identification

# KAUST 058 - Diatom of currently unknown identity





Seawater diatom isolated from North Beach at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 23°37'02.6"N 38°48'17.8"E

# Strain Identification

# KAUST 059 - Cyanobacterium of currently unknown identity





Seawater cyanobacterium isolated from North Beach at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.

Cultivation medium: Red Sea water medium with f/2 nutrients..

GPS coordinate: 23°37'02.6"N 38°48'17.8"E

## Strain Identification

# KAUST 060 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from the Dunes of Masturah desert.

Cultivation medium: BG-11.

GPS coordinate: 23°37'02.6"N 38°48'17.8"E

Strain Identification

#### **KAUST 061 - Pleodorina illinoisensis**





Freshwater *Pleodorina illinoisensis* isolated from hot springs of Wadi Zee Valley.

Cultivation medium: BG-11.


GPS coordinate: 22°43'27.6"N 39°48'37.5"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R - used in a BLAST search, the sequence identity of 99.13% shows that *Pleodorina illinoisensis* is a relative.

### KAUST 062 - Coelastrum microporum





Freshwater *Coelastrum microporum* isolated from hot springs of Wadi Zee Valley.

Cultivation medium: BG-11.

GPS coordinate: 22°43'27.6"N 39°48'37.5"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 98.24% shows that *Coelastrum microporum* is a relative.

### **KAUST 063 - Monoraphidium circinale**





Freshwater *Monoraphidium circinale* isolated from hot springs of Wadi Zee Valley.

Cultivation medium: BG-11.


GPS coordinate: 22°43'27.6"N 39°48'37.5"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase -Large subunit -RbcL F /RbcL R - used in a BLAST search, the sequence identity of 96.60% shows that *Monoraphidium circinale* is a relative.

### KAUST 064 - Chlamydomonas sp.





Freshwater "Chlamydomonas sp." isolated from hot springs of Wadi Zee Valley.

Cultivation medium: BG-11.

GPS coordinate: 22°43'27.6"N 39°48'37.5"E


### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 91.84% shows that the strain currently identified as "Chlamydomonas sp. KU108" is a relative.

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity by ITS is not conclusive. Therefore, the name is indicated in quotations.

#### **KAUST 065 – Phormidesmis molle**





Freshwater *Phormidesmis molle* isolated from hot springs of Wadi Zee Valley.

Cultivation medium: BG-11.


GPS coordinate: 22°43'27.6"N 39°48'37.5"E

# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 16S ribosomal DNA – Cya106 F/Cya781b R- used in a BLAST search, the sequence identity of 100% shows that *Phormidesmis* molle DS2\_D14 is a relative.

## KAUST 066 - Desmodesmus sp.





Freshwater *Desmodesmus* sp. isolated from hot springs of Wadi Zee Valley.

Cultivation medium: BG-11.

GPS coordinate: 22°43'27.6"N 39°48'37.5"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 100% shows that the strain currently identified as *Desmodesmus* sp. Jo\_2 is a relative.

### KAUST 067 - Scenedesmus sp.





Freshwater *Scenedesmus* sp. isolated from hot springs of Wadi Zee Valley.

Cultivation medium: BG-11.

GPS coordinate: 22°43'27.6"N 39°48'37.5"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R - used in a BLAST search, the sequence identity of 99.13% shows that *Scenedesmus* sp. KNUA019 is a relative.

\*This strain was shared with the DAB-KSA team in August/2024 for further analysis in scale-up cultivations.

### KAUST 068 - Nitzschia sp.





Seawater *Nitzschia* sp. isolated from hot springs of Wadi Zee Valley.

Cultivation medium: Red Sea water medium with f/2 nutrients..

GPS coordinate: 22°43'27.6"N 39°48'37.5"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Nitzschia* sp. is a relative.

## KAUST 069 - Anabaenopsis sp.





Freshwater *Anabaenopsis* sp. isolated from hot springs of Wadi Zee Valley.

Cultivation medium: BG-11.

GPS coordinate: 22°43'27.6"N 39°48'37.5"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 16S ribosomal DNA – Cya106 F/Cya781b R- used in a BLAST search, the sequence identity of 100% shows that *Anabaenopsis* sp. is a relative.

## KAUST 070 - Fistulifera saprophila





Seawater *Fistulifera saprophila* isolated from hot springs of Wadi Zee Valley.

Cultivation medium: Red Sea water medium with f/2 nutrients..

GPS coordinate: 22°43'27.6"N 39°48'37.5"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Fistulifera saprophila* is a relative.

## KAUST 071 – Desmodesmus sp.





Freshwater *Desmodesmus* sp. isolated from hot springs of Wadi Zee Valley.

Cultivation medium: BG-11.


GPS coordinate: 22°43'27.6"N 39°48'37.5"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 86.32% shows that *Desmodesmus* sp. is a relative.

### KAUST 072 - Chlamydomonas sp.





Freshwater "Chlamydomonas sp." isolated during a sandstorm at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 22°18'28.0"N 39°06'23.0"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 95.22% shows that the strain currently identified as "Chlamydomonas sp. YACCYB80" is a relative.

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity by ITS is not conclusive. Therefore, the name is indicated in quotations.

# KAUST 073 - Green alga of currently unknown identity





Freshwater green alga isolated during a sandstorm at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 22°18'28.0"N 39°06'23.0"E

# Strain Identification

### **KAUST 074 - Tetranephris brasiliensis**





Freshwater *Tetranephris brasiliensis* isolated during a sandstorm at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.

Cultivation medium: BG-11


GPS coordinate: 22°18'28.0"N 39°06'23.0"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 84.93% shows that *Tetranephris brasiliensis* - strain: GS3c is a relative.

# KAUST 075 - Green alga of currently unknown identity





Freshwater green alga isolated during a sandstorm at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.


Cultivation medium: BG-11

GPS coordinate: 22°18'28.0"N 39°06'23.0"E

# Strain Identification

# KAUST 076 - Green alga of currently unknown identity





Freshwater green alga isolated from Duba Corniche, at the Eiffel tower monument.

Cultivation medium: BG-11

GPS coordinate: 27°21'11.1"N 35°40'38.6"E

Strain Identification

#### **KAUST 077 - Stichococcus mirabilis**





Freshwater *Stichococcus mirabilis* isolated from Duba Corniche, at the Eiffel tower monument.

Cultivation medium: BG-11


GPS coordinate: 27°21'11.1"N 35°40'38.6"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 99.23% shows that *Stichococcus mirabilis* – strain SAG 379-3a is a relative.

### KAUST 078 - Chlamydomonas reinhardtii





Freshwater "Chlamydomonas reinhardtii" isolated from Duba Corniche, at the Eiffel tower monument.

Cultivation medium: BG-11

GPS coordinate: 27°21'11.1"N 35°40'38.6"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 91.43% shows that "Chlamydomonas reinhardtii" is a relative.

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity by ITS is not conclusive. Therefore, the name is indicated in quotations.

# KAUST 079 - Diatom of currently unknown identity





Seawater diatom isolated from Haddah Beach North, at NEOM Region.


Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 28°04'54.1"N 34°50'08.4"E

# Strain Identification

# KAUST 080 - Diatom of currently unknown identity





Seawater diatom isolated from Haddah Beach North, at NEOM Region.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 28°04'54.1"N 34°50'08.4"E

# Strain Identification

# KAUST 081 - Green alga of currently unknown identity





Seawater green alga isolated from Haddah Beach South, at NEOM Region.


Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 28°04'32.7"N 34°51'29.8"E

# Strain Identification

# KAUST 082 - Green alga of currently unknown identity





Seawater green alga isolated from Haddah Beach South, at NEOM Region.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 28°04'32.7"N 34°51'29.8"E

# Strain Identification

### KAUST 083 - Nitzschia sp.





Seawater *Nitzschia* sp. isolated from NEOM Region, at the cost of Royal Tulip Resort.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 28°02'55.6"N 35°13'00.7"E

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Nitzschia* sp. VNikulin-2024a is a relative.

# KAUST 084 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from rocks at Haddah Beach South, at NEOM Region.

Cultivation medium: BG-11

GPS coordinate: 28°04'32.7"N 34°51'29.8"E

# Strain Identification

# KAUST 085 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from rocks at Haddah Beach South, at NEOM Region.

Cultivation medium: BG-11

GPS coordinate: 28°04'32.7"N 34°51'29.8"E

Strain Identification

# KAUST 086 - Green alga of currently unknown identity





Freshwater green algae from rocks at Haddah Beach North, at NEOM Region.


Cultivation medium: BG-11

GPS coordinate: 28°04'32.7"N 34°51'29.8"E

Strain Identification

### **KAUST 087 - Chlamydomonas isabeliensis**





Freshwater "Chlamydomonas isabeliensis" isolated from rocks at Haddah Beach North, at NEOM Region.

Cultivation medium: BG-11

GPS coordinate: 28°04'32.7"N 34°51'29.8"E


## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 99.27% shows that the strain currently identified as "Chlamydomonas isabeliensis" is a relative.

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity by 18S is not conclusive. Therefore, the name is indicated in quotations.

# KAUST 088 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from rocks at Haddah Beach North, at NEOM Region.


Cultivation medium: BG-11

GPS coordinate: 28°04'32.7"N 34°51'29.8"E

Strain Identification

# KAUST 089 - Green alga of currently unknown identity





Seawater green alga isolated from rocks at Haddah Beach North, at NEOM Region.


Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 28°04'32.7"N 34°51'29.8"E

# Strain Identification

### KAUST 090 - Synechocystis sp.





Freshwater *Synechocystis* sp. isolated from the coast at Jubail.


Cultivation medium: BG-11


GPS coordinate: 27°05'02.0"N 49°32'07.9"E

## Strain Identification

Besides the morphological taxonomy, taxonomic classification was evaluated by first performing a genome assembly of the strain, followed by 16s rRNA sequence isolation using Barrnap and a k-mer based classification using Kraken2. rRNA sequences were searched against the NCBI database using BLAST. Both analyses show "Synechocystis" sp. PCC 6803" is a relative, with a BLAST results showing a sequence identity of 99.60%.

# KAUST 091 - Green alga of currently unknown identity






Seawater green alga isolated from the coast at Jubail. Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 27°05'02.0"N 49°32'07.9"E

# Strain Identification

# KAUST 092 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from a bridge close to the coast at Jubail.


Cultivation medium: BG-11

GPS coordinate: 27°07'14.0"N 49°34'16.0"E

# Strain Identification

# KAUST 093 – Cyanobacterium of currently unknown identity





Seawater cyanobacterium isolated from a bridge close to the coast at Jubail.


Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 27°07'14.0"N 49°34'16.0"E

# Strain Identification

#### **KAUST 094 - Heterochlorella luteoviridis**





Freshwater *Heterochlorella luteoviridis* isolated from a bridge close to the coast at Jubail.

Cultivation medium: BG-11

GPS coordinate: 27°07'14.0"N 49°34'16.0"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase-Large subunit-RbcL F/RbcL R - used in a BLAST search, the sequence identity of 95.83% shows that *Heterochlorella luteoviridis* is a relative.

## KAUST 095 - Chlamydomonas sp.





Freshwater "Chlamydomonas sp." isolated from Jubail area.

Cultivation medium: BG-11

GPS coordinate: 27°05'02.0"N 49°32'22.3"E

## Strain Identification

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity by 18S is not conclusive.

Therefore, the name is indicated in quotations.

# KAUST 096 - Green alga of currently unknown identity





Freshwater green alga isolated from Jubail area.

Cultivation medium: BG-11

GPS coordinate: 27°05'02.0"N 49°32'22.3"E

# Strain Identification

# KAUST 097 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from Jubail area.


Cultivation medium: BG-11

GPS coordinate: 27°05'02.0"N 49°32'22.3"E

Strain Identification

# KAUST 098 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from Jubail area.

Cultivation medium: BG-11

GPS coordinate: 27°05'02.0"N 49°32'22.3"E

Strain Identification

# KAUST 099 - Green alga of currently unknown identity





Freshwater green alga isolated from Mangroves at Dammam area (Abu Hadriah Highway).

Cultivation medium: BG-11

GPS coordinate: 26°49'26.0"N 49°34'37.0"E

# Strain Identification

### **KAUST 100 - Viridiparva madeirensis**





Freshwater *Viridiparva madeirensis* isolated from a bark tree at Deffi Park at Jubail.

Cultivation medium: BG-11

GPS coordinate: 27°08'43.0"N 49°32'07.0"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase -Large subunit -RbcL F /RbcL R - used in a BLAST search, the sequence identity of 95.41% shows that *Viridiparva madeirensis* is a relative.

# KAUST 101 - Green alga of currently unknown identity





Freshwater green alga isolated from a bark tree at Deffi Park at Jubail.

Cultivation medium: BG-11

GPS coordinate: 27°08'43.0"N 49°32'07.0"E

# Strain Identification

#### KAUST 102 - Stichococcus sp.





Freshwater *Stichococcus sp.* isolated from a bark tree at Deffi Park at Jubail.

Cultivation medium: BG-11

GPS coordinate: 27°08'43.0"N 49°32'07.0"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase-Large subunit-RbcL F/RbcL R - used in a BLAST search, the sequence identity of 97.73% shows that *Stichococcus sp.* is a relative.

# KAUST 103 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from the Deserts regions of Al Ula, located in the Medina Region in north-western Saudi Arabia.

Cultivation medium: BG-11

GPS coordinate: 26°36'59.99"N 37°54'59.99"E

## Strain Identification

# KAUST 104 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from the Deserts regions of Al Ula, located in the Medina Region in north-western Saudi Arabia.

Cultivation medium: BG-11

GPS coordinate: 26°36'59.99"N 37°54'59.99"E

## Strain Identification

# KAUST 105 - Green alga of currently unknown identity





Freshwater green alga isolated from the area of Jeddah Corniche.


Cultivation medium: BG-11

GPS coordinate: 21°35'43.8"N 39°06'19.8"E

# Strain Identification

# KAUST 106 - Green alga of currently unknown identity





Freshwater green alga isolated from the area of Jeddah Corniche.


Cultivation medium: BG-11

GPS coordinate: 21°35'43.8"N 39°06'19.8"E

# Strain Identification

# KAUST 107 - Green alga of currently unknown identity





Freshwater green alga isolated from the area of Jeddah Corniche.


Cultivation medium: BG-11

GPS coordinate: 21°35'43.8"N 39°06'19.8"E

# Strain Identification

# KAUST 108 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from the area of Jeddah Corniche.

Cultivation medium: BG-11

GPS coordinate: 21°35'43.8"N 39°06'19.8"E

# Strain Identification

# KAUST 109 - Green alga of currently unknown identity





Freshwater green alga isolated from the area of Khulays, Makkah Region.

Cultivation medium: BG-11

GPS coordinate: 22°8'39"N 39°35'49"E

# Strain Identification

KAUST 110 - Green alga of currently unknown identity





Freshwater green alga isolated from the Deserts regions of Al Ula, located in the Medina Region in north-western Saudi Arabia.

Cultivation medium: BG-11

GPS coordinate: 26°36'59.99"N 37°54'59.99"E

# Strain Identification

#### KAUST 111 – Chlorella sorokiniana



Freshwater *Chlorella sorokiniana* isolated from the wastewater treatment plants in Makkah by Mohammad Khalil Monjed's team.

Cultivation medium: BG-11

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 99.55% shows that *Chlorella sorokiniana* is a relative.

#### KAUST 112 – Parachlorella kessleri



Freshwater *Parachlorella kessleri* isolated from the wastewater treatment plants in Makkah by Mohammad Khalil Monjed's team.

Cultivation medium: BG-11

# Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Parachlorella kessleri* is a relative.

### **KAUST 113** – *Chlorella vulgaris*



Freshwater *Chlorella vulgaris* isolated from the wastewater treatment plants in Makkah by Mohammad Khalil Monjed's team.

Cultivation medium: BG-11

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Chlorella vulgaris* is a relative.

#### KAUST 114 - Chlorella sp.




Freshwater *Chlorella* sp. isolated from the wastewater treatment plants in Makkah by Mohammad Khalil Monjed's team.

Cultivation medium: BG-11

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Chlorella* sp. YACCYB105 is a relative.

#### KAUST 115 - Chlorella sorokiniana



Seawater *Chlorella sorokiniana* isolated from non-treated wastewater at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°44'03.0"N 39°46'34.0"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Chlorella sorokiniana* isolate HH01. BZ-2019 is a relative.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

## KAUST 116 - Micractinium sp.



Freshwater *Micractinium* sp. isolated from treated wastewater at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: BG-11


GPS coordinate: 22°44'03.0"N 39°46'34.0"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Micractinium* sp. ACSSI 332 is a relative.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

## KAUST 117 - Chlorella sp.



Freshwater *Chlorella* sp. isolated from non-treated wastewater at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: BG-11

GPS coordinate: 22°44'03.0"N 39°46'34.0"E


#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R- used in a BLAST search, the sequence identity of 100% shows that *Chlorella* sp. KMMCC 1468 is a relative.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

### KAUST 118 – Leptolyngbya breviarticulata





Seawater *Leptolyngbya breviarticulata* isolated from a mud sample from Site 48 in the Empty Quarter Project.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 20°34'07.1"N 53°05'38.6"E

### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 16S ribosomal DNA – Cya106 F/Cya781b R- used in a BLAST search, the sequence identity of 99.64% shows that *Leptolyngbya breviarticulata ISB 46* is a relative.

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

### KAUST 119 - Green alga of currently unknown identity



Seawater green alga isolated from solid biofilm at one of the farms at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°3'7"N 39°8'34"E

### Strain Identification

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

### KAUST 120 - Cyanobacterium of currently unknown identity



Freshwater cyanobacterium isolated from a biofilm outside pipes at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: BG-11

GPS coordinate: 22°44'03.0"N 39°46'34.0"E

### Strain Identification

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

## **KAUST 121** – Diatiom of currently unknown identity





Seawater diatom isolated from a water and clay sample from Site 59 in the Empty Quarter Project.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 20°31'24.7"N 54°47'13.7"

### Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

## KAUST 122 - Green alga of currently unknown identity





Freshwater green alga isolated from Site 54 in the Empty Quarter Project.

Cultivation medium: BG-11


GPS coordinate: 20°55'24"N 53°57'43.936"

## Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

### KAUST 123 - Cyanobacterium of currently unknown identity





Seawater cyanobacterium isolated from a water and clay sample from Site 60 in the Empty Quarter Project.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 20°42'22.948"N 54°43'55.58"E

### Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

#### **KAUST 124** – Diatom of currently unknown identity



Seawater diatom isolated from non-treated wastewater at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°44'03.0"N 39°46'34.0"

### Strain Identification

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

### KAUST 125 - Diatom of currently unknown identity



Seawater diatom isolated from a biofilm outside pipes at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°44'03.0"N 39°46'34.0"E

### Strain Identification

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

### KAUST 126 - Cyanobacterium of currently unknown identity





Seawater cyanobacterium isolated from a water and clay sample from Site 60 in the Empty Quarter Project.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 20°42'22.948"N 54°43'55.58"E

### Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

### **KAUST 127** – Cyanobacterium of currently unknown identity





Seawater cyanobacterium isolated from a water and clay sample from Site 59 in the Empty Quarter Project.

Cultivation medium: Red Sea water medium with f/2 nutrients.


GPS coordinate: 20°31'24.7"N 54°47'13.7"

### Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

## KAUST 128 - Green alga of currently unknown identity





Freshwater green alga isolated from Site 54 in the Empty Quarter Project.

Cultivation medium: BG-11

GPS coordinate: 20°55'24"N 53°57'43.936"

## Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

#### **KAUST 129** – Diatom of currently unknown identity



Seawater diatom isolated from solid biofilm at one of the farms at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.


Cultivation medium: BG-11

GPS coordinate: 22°3'7"N 39°8'34"E

#### Strain Identification

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

#### KAUST 130 - Asterarcys sp.



Freshwater *Asterarcys* sp. isolated from solid biofilm at one of the farms at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: BG-11

GPS coordinate: 22°3'7"N 39°8'34"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase-Large subunit-RbcL F/RbcL R - used in a BLAST search, the sequence identity of 97.64% shows that *Asterarcys* sp. GA4 is a relative.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

#### KAUST 131 - Chlamydomonas sp.



Freshwater "Chlamydomonas sp". isolated from one of the farms at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: BG-11

GPS coordinate: 22°3'7"N 39°8'34"E

#### Strain Identification


Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of rbcL\_F1 NC\_005353.1:123917/ rbcL\_R8 NC\_005353.1:122737- used in a BLAST search, the sequence identity of 99.53% shows that "Chlamydomonas sp." HC1 is a relative.

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity is not conclusive.

Therefore, the name is indicated in quotations.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

#### KAUST 132 - Chlorella sp.




Seawater *Chlorella* sp. isolated from Red Sea water samples at Jeddah Corniche by Mohammad Sherif Edris's team.

Cultivation medium: Red Sea water medium with f/2 nutrients.

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R - used in a BLAST search, the sequence identity of 100% shows that *Chlorella* sp. KMMCC C-66 is a relative.

#### KAUST 133 – *Tetraselmis* sp.




Seawater *Tetraselmis* sp. isolated from Red Sea water samples at Jeddah Corniche by Mohammad Sherif Edris's team.

Cultivation medium: Red Sea water medium with f/2 nutrients.

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase-Large subunit-RbcL F/RbcL R - used in a BLAST search, the sequence identity of 99.75% shows that *Tetraselmis* sp. KMMCC 84 is a relative.

#### KAUST 134 – Chlorella sorokinina




Seawater *Chlorella sorokiniana* isolated from Red Sea water samples at Jeddah Corniche by Mohammad Sherif Edris's team. Cultivation medium: Red Sea water medium with f/2 nutrients.

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Ribulose-1,5-bisphosphate carboxylase/oxygenase-Large subunit-RbcL F/RbcL R - used in a BLAST search, the sequence identity of 100% shows that *Chlorella sorokiniana* isolate 275\_4 is a relative.

#### KAUST 135 – Dunaliella tertiolecta




Seawater *Dunaliella tertiolecta* isolated from Red Sea water samples at Jeddah Corniche by Mohammad Sherif Edris's team. Cultivation medium: Red Sea water medium with f/2 nutrients.

## Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R - used in a BLAST search, the sequence identity of 100% shows that *Dunaliella tertiolecta* straisn CCAP 19/7C is a relative.

#### KAUST 136 - Monoraphidium subclavatum



Freshwater *Monoraphidium subclavatum* isolated from solid biofilm at one of the farms at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: BG-11

GPS coordinate: 22°3'7"N 39°8'34"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of Internal transcribed spacer 1 - ITS1 F (modified)/ITS2T R - used in a BLAST search, the sequence identity of 98.39% shows that *Monoraphidium subclavatum* strain Ga035 is a relative.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

#### **KAUST 137** – Diatom of currently unknown identity



Seawater diatom isolated at Saudi Radwa Food Company
Limited facilities, located in Al-Bi'ar in the Makkah Province.


Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°44'03.0"N 39°46'34.0"E

#### Strain Identification

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

#### KAUST 138 - Chlamydomonas sp.



Freshwater "Chlamydomonas sp." isolated from one of the farms at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: BG-11

GPS coordinate: 22°3'7"N 39°8'34"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R - used in a BLAST search, the sequence identity of 100% shows that "Chlamydomonas sp." YACCYB5is a relative.

\*Chlamydomonas taxonomy is currently undergoing revision, and true identity by 18S ribosomal DNA – Cdm F/Cdm R is not conclusive. Therefore, the name is indicated in quotations.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

## KAUST 139 – *Porphyridium* sp.





Seawater *Porphyridium* sp. isolated from North Beach at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.


Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 23°37'02.6"N 38°48'17.8"E

## Strain Identification

## KAUST 140 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from Site HQ25-KY7 in the Empty Quarter.

Cultivation medium: BG-11


GPS coordinate: 19°39'5.202"N 57°25'0.552"E

## Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Volker Vahrenkamp's team.

#### **KAUST 141** – Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from a water and clay sample from Site 60 in the Empty Quarter Project.

Cultivation medium: BG-11

GPS coordinate: 20°42'22.948"N 54°43'55.58"E

#### Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

## KAUST 142 - Green alga of currently unknown identity





Seawater green alga isolated from Site 54 in the Empty Quarter Project.

Cultivation medium: Red Sea water medium with f/2 nutrients.


GPS coordinate: 20°55'24"N 53°57'43.936"

## Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

## **KAUST 143** – Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from a water and clay sample from Site 55 in the Empty Quarter Project.

Cultivation medium: BG-11


GPS coordinate: 20°54'8.181"N 54°0'8.77536"E

#### Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

## KAUST 144 – Cyanobacterium of currently unknown identity





Seawater cyanobacterium isolated from Site 54 in the Empty Quarter Project.

Cultivation medium: Red Sea water medium with f/2 nutrients.


GPS coordinate: 20°55'24"N 53°57'43.936"

## Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

## KAUST 145 – Cyanobacterium of currently unknown identity





Seawater cyanobacterium isolated from Site 54 in the Empty Quarter Project.

Cultivation medium: Red Sea water medium with f/2 nutrients.


GPS coordinate: 20°55'24"N 53°57'43.936"

## Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

#### KAUST 146 – Cyanobacterium of currently unknown identity





Seawater cyanobacterium isolated from a water and clay sample from Site 60 in the Empty Quarter Project.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 20°42'22.948"N 54°43'55.58"E

#### Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Robert Hoehndorf's team, under the Empty Quarter Project.

#### KAUST 147 – Scenedesmus sp.



Seawater *Scenedesmus* sp. isolated from solid biofilm at one of the farms at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°3'7"N 39°8'34"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R - used in a BLAST search, the sequence identity of 100% shows that 100% *Scenedesmus* sp. ISC 121 is a relative.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

#### KAUST 148 - Synechocystis sp.



Freshwater *Synechocystis* sp. isolated from solid biofilm at one of the farms at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°3'7"N 39°8'34"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 16S ribosomal DNA – Cya106 F/Cya781b R- used in a BLAST search, the sequence identity of 100% shows that 100% *Synechocystis* sp. PCC 6803 is a relative.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

#### KAUST 149 – Rhizoclonium sp.



Seawater *Rhizoclonium* sp. isolated from a biofilm outside pipes at Saudi Radwa Food Company Limited facilities, located in Al-Bi'ar in the Makkah Province.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°44'03.0"N 39°46'34.0"E

#### Strain Identification

Besides the morphological taxonomy, according to the DNA amplification for the partial sequence of 18S ribosomal DNA – Cdm F/Cdm R - used in a BLAST search, the sequence identity of 100% shows that *Rhizoclonium* sp. is a relative.

\*This strain resulted from environmental samples taken on a field trip to Saudi Radwa Food Company Limited facilities together with Development of Algal Biotechnology in the Kingdom of Saudi Arabia (DAB-KSA) team, KAUST Beacon Development, a project funded by the Ministry of Environment Water & Agriculture (MEWA). The visit was organized through the KAUST Center of Excellence in Sustainable Food Security.

## KAUST 150 – Cyanobacterium of currently unknown identity





 $Freshwater\ cyanobacterium\ isolated\ from\ Site$ 

HQ25-KY10 in the Empty.

Cultivation medium: BG-11

GPS coordinate: 20°58'26.952"N 58°13'15.018"E

## Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Volker Vahrenkamp's team.

## KAUST 151 – Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from Site HQ25-KY06 in the Empty Quarter.

Cultivation medium: BG-11

GPS coordinate: 19°41'38.01"N 57°23'53.628"E

## Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Volker Vahrenkamp's team.

#### KAUST 152 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated from Site HQ25-KY06 in the Empty Quarter.

Cultivation medium: BG-11

GPS coordinate: 19°41'38.01"N 57°23'53.628"E

#### Strain Identification

\*This strain resulted from environmental samples taken on a field trip by Volker Vahrenkamp's team.

## KAUST 153 - Cyanobacterium of currently unknown identity





Freshwater cyanobacterium isolated during a sandstorm at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.

Cultivation medium: BG-11

GPS coordinate: 22°18'28.0"N 39°06'23.0"E

#### Strain Identification

## KAUST 154 - Diatom of currently unknown identity





Seawater diatom isolated from North Beach at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.


Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 23°37'02.6"N 38°48'17.8"E

## Strain Identification

## KAUST 155 - Green alga of currently unknown identity






Freshwater green alga isolated during a sandstorm at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region. Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 22°18'28.0"N 39°06'23.0"E

#### Strain Identification

## KAUST 156 - Green alga of currently unknown identity





Seawater green alga isolated from North Beach at King Abdullah University of Science and Technology (Thuwal, Makkah), in the mid-Red Sea coast region.

Cultivation medium: Red Sea water medium with f/2 nutrients.

GPS coordinate: 23°37'02.6"N 38°48'17.8"E

## Strain Identification

# KSA Algae Living Library

Microalgae and cyanobacteria isolated from the Kingdom of Saudi Arabia

# **General inquiries**

Prof. Kyle J. Lauersen

kyle.lauersen@kaust.edu.sa

Dr. Bárbara Bastos de Freitas

barbara.bastosdefreitas@kaust.edu.sa

